

Ref : FT.L0096an
Version date : 04/12/13

DMEM Ham's F12
w/o L-Glutamine, w/ 25mM Hepes

CAT N° : SLD-521-500

Theoretical pH : 7.2 ± 0.3

Osmolality : 304 mOsm/kg $\pm 10\%$

Colour : Red solution

Storage conditions : +2°C to +8°C in the dark

Shelf life : 24 months

Sterility tests :

- Bacteria in aerobic and anaerobic conditions
- Fungi and yeasts

Endotoxin : <1 EU/ml

Cell growth test :

Medium tested for the ability to support cell growth with Hela line.

Composition : Displayed on web site; also available on request

Recommended use :

- Respect storage conditions of the product
- Do not use the product after its expiry date
- Store product in an area protected from light (not necessary for saline solutions).
- Manipulate the product in aseptic conditions (e.g. : under laminar air flow)
- Wear clothes adapted to the manipulation of the product to avoid contamination (e.g. : gloves, mask, hygiene cap, overall...)

The product is intended to be used in vitro, in laboratory only. Do not use it in therapy, human or veterinary applications.

Description :

Studies to determine the nutritional requirements of many cells have been in progress since Eagle's first reports. The major essential nutrients were identified and work became focused on the media requirements of individual cell types. Many media designed for these purposes are now available. Among the first of these media, developed initially to study hormonal requirements of cells in culture, was a mixture of DMEM medium and Ham's F12 medium, known as DMEM Ham's F12.

Uses :

Supplements, such as antibiotics, should be added as sterile supplements to the medium. Storage conditions and shelf-life of supplemented products will be affected by the nature of the supplements. Add 12.5 ml/l of L-Glutamine 100X, 200mM (CAT N° : SRL-810-500) or 365 mg/l of L-Glutamine (CAT N° : SPL-134-1KG) before using this medium.

Signs of deterioration :

Medium should be clear and free of particulate and flocculent material.

Do not use this medium if it is cloudy or contains precipitate.

Other evidence of deterioration may include colour change or degradation of physical or performance characteristics.